



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 894 854 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
03.02.1999 Bulletin 1999/05

(21) Application number: 97309791.8

(22) Date of filing: 04.12.1997

(51) Int. Cl.⁶: C12N 15/12, C07K 14/705,
A61K 38/17, C12Q 1/68,
G01N 33/68, G01N 33/50,
C07K 16/28

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 29.07.1997 US 902294

(71) Applicant:
SMITHKLINE BEECHAM CORPORATION
Philadelphia Pennsylvania 19103 (US)

(72) Inventor: Zhu, Yuan
King of Prussia, Pennsylvania 19406 (US)

(74) Representative:
Crump, Julian Richard John et al
FJ Cleveland,
40-43 Chancery Lane
London WC2A 1JQ (GB)

(54) **A novel splicing variant of the epstein-barr virus-induced g-protein coupled receptor**

(57) EBI 3 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing EBI 3 polypeptides and polynucleotides in the design of protocols for the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others and diagnostic assays for such conditions.

EP 0 894 854 A2

Description

FIELD OF INVENTION

- 5 [0001] This invention relates to newly identified polynucleotides, polypeptides encoded by them and to the use of such polynucleotides and polypeptides, and to their production. More particularly, the polynucleotides and polypeptides of the present invention relate to G-protein coupled receptor family, hereinafter referred to as EBI 3. The invention also relates to inhibiting or activating the action of such polynucleotides and polypeptides.

10 BACKGROUND OF THE INVENTION

- [0002] It is well established that many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers, e.g., cAMP (Lefkowitz, Nature, 1991, 351:353-354). Herein these proteins are referred to as proteins participating in pathways with G-proteins or PPG
15 proteins. Some examples of these proteins include the GPC receptors, such as those for adrenergic agents and dopamine (Kobilka, B.K., et al., Proc. Natl Acad. Sci., USA, 1987, 84:46-50; Kobilka, B.K., et al., Science, 1987, 238:650-656; Bunzow, J.R., et al., Nature, 1988, 336:783-787), G-proteins themselves, effector proteins, e.g., phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins, e.g., protein kinase A and protein kinase C (Simon, M.I., et al., Science, 1991, 252:802-8).

- 20 [0003] For example, in one form of signal transduction, the effect of hormone binding is activation of the enzyme, adenylyl cyclase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide GTP. GTP also influences hormone binding. A G-protein connects the hormone receptor to adenylyl cyclase. G-protein was shown to exchange GTP for bound GDP when activated by a hormone receptor. The GTP-carrying form then binds to activated adenylyl cyclase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its
25 basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

- [0004] The membrane protein gene superfamily of G-protein coupled receptors has been characterized as having seven putative transmembrane domains. The domains are believed to represent transmembrane α -helices connected by extracellular or cytoplasmic loops. G-protein coupled receptors include a wide range of biologically active receptors,
30 such as hormone, viral, growth factor and neuroreceptors.

- [0005] G-protein coupled receptors (otherwise known as 7TM receptors) have been characterized as including these seven conserved hydrophobic stretches of about 20 to 30 amino acids, connecting at least eight divergent hydrophilic loops. The G-protein family of coupled receptors includes dopamine receptors which bind to neuroleptic drugs used for treating psychotic and neurological disorders. Other examples of members of this family include, but are not limited to,
35 calcitonin, adrenergic, endothelin, cAMP, adenosine, muscarinic, acetylcholine, serotonin, histamine, thrombin, kinin, follicle stimulating hormone, opsins, endothelial differentiation gene-1, rhodopsins, odorant, and cytomegalovirus receptors.

- [0006] Most G-protein coupled receptors have single conserved cysteine residues in each of the first two extracellular loops which form disulfide bonds that are believed to stabilize functional protein structure. The 7 transmembrane
40 regions are designated as TM1, TM2, TM3, TM4, TM5, TM6, and TM7. TM3 has been implicated in signal transduction.

- [0007] Phosphorylation and lipidation (palmitoylation or farnesylation) of cysteine residues can influence signal transduction of some G-protein coupled receptors. Most G-protein coupled receptors contain potential phosphorylation sites within the third cytoplasmic loop and/or the carboxy terminus. For several G-protein coupled receptors, such as the β -adrenoreceptor, phosphorylation by protein kinase A and/or specific receptor kinases mediates receptor desensitiza-
45 tion.

- [0008] For some receptors, the ligand binding sites of G-protein coupled receptors are believed to comprise hydrophilic sockets formed by several G-protein coupled receptor transmembrane domains, said sockets being surrounded by hydrophobic residues of the G-protein coupled receptors. The hydrophilic side of each G-protein coupled receptor transmembrane helix is postulated to face inward and form polar a ligand binding site. TM3 has been impli-
50 cated in several G-protein coupled receptors as having a ligand binding site, such as the TM3 aspartate residue. TM5 serines, a TM6 asparagine and TM6 or TM7 phenylalanines or tyrosines are also implicated in ligand binding.

- [0009] G-protein coupled receptors can be intracellularly coupled by heterotrimeric G-proteins to various intracellular enzymes, ion channels and transporters (see, Johnson et al., Endoc. Rev., 1989, 10:317-331). Different G-protein α -subunits preferentially stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation
55 of cytoplasmic residues of G-protein coupled receptors has been identified as an important mechanism for the regulation of G-protein coupling of some G-protein coupled receptors. G-protein coupled receptors are found in numerous sites within a mammalian host.

- [0010] Over the past 15 years, nearly 350 therapeutic agents targeting 7 transmembrane (7 TM) receptors have been

successfully introduced onto the market.

[0011] This indicates that these receptors have an established, proven history as therapeutic targets. Clearly there is a need for identification and characterization of further receptors which can play a role in preventing, ameliorating or correcting dysfunctions or diseases, including, but not limited to, infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome.

SUMMARY OF THE INVENTION

[0012] In one aspect, the invention relates to EBI 3 polypeptides and recombinant materials and methods for their production. Another aspect of the invention relates to methods for using such EBI 3 polypeptides and polynucleotides. Such uses include the treatment of infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others. In still another aspect, the invention relates to methods to identify agonists and antagonists using the materials provided by the invention, and treating conditions associated with EBI 3 imbalance with the identified compounds. Yet another aspect of the invention relates to diagnostic assays for detecting diseases associated with inappropriate EBI 3 activity or levels.

DESCRIPTION OF THE INVENTION

Definitions

[0013] The following definitions are provided to facilitate understanding of certain terms used frequently herein.

[0014] "EBI 3" refers, among others, to a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2, or an allelic variant thereof.

[0015] "Receptor Activity" or "Biological Activity of the Receptor" refers to the metabolic or physiologic function of said EBI 3 including similar activities or improved activities or these activities with decreased undesirable side-effects. Also included are antigenic and immunogenic activities of said EBI 3.

[0016] "EBI 3 gene" refers to a polynucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 1 or allelic variants thereof and/or their complements.

[0017] "Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

[0018] "Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

[0019] "Polynucleotide" generally refers to any polyribonucleotide or polydeoxiribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications has been made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

[0020] "Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides

may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Posttranslational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62.

[0021] "Variant" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

[0022] "Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques. See, e.g.: (*COMPUTATIONAL MOLECULAR BIOLOGY*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS*, Smith, D.W., ed., Academic Press, New York, 1993; *COMPUTER ANALYSIS OF SEQUENCE DATA, PART I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY*, von Heinje, G., Academic Press, 1987; and *SEQUENCE ANALYSIS PRIMER*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in *Guide to Huge Computers*, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipton, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCS program package (Devereux, J., *et al.*, *Nucleic Acids Research* (1984) 12(1):387), BLASTP, BLASTN, FASTA (Atschul, S.F. *et al.*, *J Molec Biol* (1990) 215:403).

[0023] As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence of SEQ ID NO: 1 is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence of SEQ ID NO: 1. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5 or 3 terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in

one or more contiguous groups within the reference sequence.

[0024] Similarly, by a polypeptide having an amino acid sequence having at least, for example, 95% identity to a reference amino acid sequence of SEQ ID NO:2 is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of SEQ ID NO: 2. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

Polypeptides of the Invention

[0025] In one aspect, the present invention relates to EBI 3 polypeptides (or EBI 3 proteins). The EBI 3 polypeptides include the polypeptide of SEQ ID NO:2; as well as polypeptides comprising the amino acid sequence of SEQ ID NO:2; and polypeptides comprising the amino acid sequence which have at least 80% identity to that of SEQ ID NO:2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Also included within EBI 3 polypeptides are polypeptides having the amino acid sequence which have at least 80% identity to the polypeptide having the amino acid sequence of SEQ ID NO: 2 over its entire length, and still more preferably at least 90% identity, and even still more preferably at least 95% identity to SEQ ID NO: 2. Furthermore, those with at least 97-99% are highly preferred. Preferably EBI 3 polypeptides exhibit at least one biological activity of the receptor.

[0026] The EBI 3 polypeptides may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[0027] Fragments of the EBI 3 polypeptides are also included in the invention. A fragment is a polypeptide having an amino acid sequence that entirely is the same as part, but not all, of the amino acid sequence of the aforementioned EBI 3 polypeptides. As with EBI 3 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101 to the end of EBI 3 polypeptide. In this context "about" includes the particularly recited ranges larger or smaller by several, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes.

[0028] Preferred fragments include, for example, truncation polypeptides having the amino acid sequence of EBI 3 polypeptides, except for deletion of a continuous series of residues that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Other preferred fragments are biologically active fragments. Biologically active fragments are those that mediate receptor activity, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also included are those that are antigenic or immunogenic in an animal, especially in a human.

[0029] Preferably, all of these polypeptide fragments retain the biological activity of the receptor, including antigenic activity. Variants of the defined sequence and fragments also form part of the present invention. Preferred variants are those that vary from the referents by conservative amino acid substitutions – i.e., those that substitute a residue with another of like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination.

[0030] The EBI 3 polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

Polynucleotides of the Invention

[0031] Another aspect of the invention relates to EBI 3 polynucleotides. EBI 3 polynucleotides include isolated polynucleotides which encode the EBI 3 polypeptides and fragments, and polynucleotides closely related thereto. More specifically, EBI 3 polynucleotide of the invention include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO: 1 encoding a EBI 3 polypeptide of SEQ ID NO: 2, and polynucleotide having the particular sequence of SEQ ID NO:1. EBI 3 polynucleotides further include a polynucleotide comprising a nucleotide sequence that has at least 80% identity over its entire length to a nucleotide sequence encoding the EBI 3 polypeptide of SEQ ID NO:2, and a polynucleotide comprising a nucleotide sequence that is at least 80% identical to that of SEQ ID NO: 1 over its entire length. In this regard, polynucleotides at least 90% identical are particularly preferred, and those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred and those with at least 98-99% are most highly preferred, with at least 99% being the most preferred. Also included under EBI 3 polynucleotides are a nucleotide sequence which has sufficient identity to a nucleotide sequence contained in SEQ ID NO: 1 to hybridize under conditions useable for amplification or for use as a probe or marker. The invention also provides polynucleotides which are complementary to such EBI 3 polynucleotides.

[0032] EBI 3 of the invention is structurally related to other proteins of the G-protein coupled receptor family, as shown by the results of sequencing the cDNA of Table 1 (SEQ ID NO: 1) encoding human EBI 3. The cDNA sequence of SEQ ID NO: 1 contains an open reading frame (nucleotide number 4 to 1087) encoding a polypeptide of 361 amino acids SEQ ID NO:2. The amino acid sequence of Table 2 (SEQ ID NO:2) has about 94.5% identity (using Lipman-Pearson Protein Alignment) in 361 amino acid residues with human EBI 1 (M. Birkenbach et al., J. Virol. 67:2209-2220, 1993). In fact, EBI 3 is a splicing variant of EBI 1, missing 17-amino acids between TM2 and TM3. The nucleotide sequence of Table 1 (SEQ ID NO: 1) has about 92.8% identity (using Martinez/Needleman-Wunsch DNA Alignment) in 1094 nucleotide residues with human EBI 1 (M. Birkenbach et al., J. Virol. 67:2209-2220, 1993). In fact, EBI 3 has an internal in frame deletion which contains 50 nucleotide residues between TM2 and TM3. Thus EBI 3 polypeptides and polynucleotides of the present invention are expected to have, inter alia, similar biological functions/properties to their homologous polypeptides and polynucleotides, and their utility is obvious to anyone skilled in the art.

Table 1^a

30	ACCATGGATT TGGGTAAACCAATGAAAAGCGT GCTGGTGG 40
	TGGCTCT CCTTGT CATT TT CCAGGTATG CCTGTGT CAAgA 80
	TGAGGT CACGGACgATTACAT CGGAgACAACACCACAGTG 120
35	GACTACACTTTGTT CGAgTCTTTGTGCT CCAAgAAGGACG 160
	TGCGGAAC TTTAAAGCCTGGTT CCTCCCTATCATGTACTC 200
	CATCATTTGTTT CGTGGGCCTACTGGGCAATGGGCTGGT C 240
	GTGTTGACCTATATCTATTTCAAgAGGCTCAAgACCATGA 280
40	COGATACCTACCTGCTCAACCTGGGGTGGCAGACATCCT 320
	CTTCTCCTGACCCTTCCCTTCTGGGCCTACAGCGGGCC 360
	AAGTCCTGGGTCTTCAGTGGCATGCTCCTACTTCTTTGCA 400
45	TCAGCATTGACGCTACGTGGCCATOGTCCAGGCTGTCTC 440
	AGCTCACOGCCACOGTGCCOGGTCTTCTCATCAGCAAG 480
	CTGTCTGTGTGGGCATCTGGATACTAGCCACAGTGCTCT 520
50	CCATCCCAGAGCTCCTGTACAGTGACCTCCAGAGGAGCAG 560
	CAGTGAGCAAGCGATGCGATGCTCTCTCATCACAGAGCAT 600
	GTGGAGGCCTTTATCACCATCCAGGTGGCCCAgaTGGTGA 640
55	TCGGCTTTCTGGTCCCCCTGCTGGCCATGAGCTTCTGTTA 680
	CCTTGT CATCATCOGCACCcTGcTCCAGGCAOGCAAcTTT 720

5 GAG CG CAACAAGGCCAT CAAGGTGAT CAT OGCTGTGGT CG 760
 TGGT CTT CAT AGT CTT CCAGCTGCCcTACAATGGGGTGGT 800
 CCTGGCCCAGACGGTGGCCAACCTT CAACAT CACCAGTAGC 840
 ACCTGTGAGCT CAGT AAG CAACT CAACAT OGCTACGACG 880
 10 T CACCTACAGCCTGGCCTGCGT CCGCTGCTGCGT CAACCC 920
 TTT CTTGTACGCCTT CAT CGGCGT CAAGTT CCGCAACGAT 960
 CT CTT CAAGCT CTT CAAGGACCTGGGCTGCCT CAGCCAGG 1000
 AGCAGCT CCGGCAGTGGT CTT CCTGT OGGCACAT CCGGCG 1040
 15 CT CCT CCATGAGTGTGGAGGCOGAGACCACCACCACCTTC 1080
 TCCCEATAAT CTAG 1094

^a A nucleotide sequence of a human EBI 3 (SEQ ID NO: 1).

Table 2^b

25 MDLGKPMK SVLVVALLVI FQVCLQDEVTDDYIGDNTTVD 40
 YTLFE SLCSKKDV RNFAWFLPIMY SII CFVGLLGNGLVV 80
 LTYIYFK RLKTM TD TYLLNLAVAD ILFLLTLPFWAY SAAK 120
 30 SWVFSGMLLLLCI SID RYVAIVQAV SAH RH RARVLLI SKL 160
 SCVGIWILATVLSIPELLYSDLQRSSEQAMRCSLITEHV 200
 EAFITIQVQMVI GFLVPLLAMSF CYLVII RTLLQARNFE 240
 35 RNKAIKVIIAVVVVFIVFQLPYNGVVLAQTVANFNIT SST 280
 CELSKQLNIAYDVTY SLACV RCCVNF FLYAFIGVKFRNDL 320
 FKLFDLGCLSQEQLRQWSSCRH IRRSSMSVEAETTTTFS 360
 40 P 362

^b An amino acid sequence of a human EBI 3 (SEQ ID NO:2).

45

[0033] One polynucleotide of the present invention encoding EBI 3 may be obtained using standard cloning and screening, from a cDNA library derived from mRNA in cells of human spleen using the expressed sequence tag (EST) analysis (Adams, M.D., *et al. Science* (1991) 252:1651-1656; Adams, M.D. *et al., Nature*, (1992) 355:632-634; Adams, M.D., *et al., Nature* (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

[0034] The nucleotide sequence encoding EBI 3 polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in Table 1 (nucleotide number 4 to 1087 of SEQ ID NO:1), or it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2.

55 [0035] When the polynucleotides of the invention are used for the recombinant production of EBI 3 polypeptide, the polynucleotide may include the coding sequence for the mature polypeptide or a fragment thereof, by itself; the coding sequence for the mature polypeptide or fragment in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro or prepro- protein sequence, or other fusion peptide portions. For

example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

[0036] Further preferred embodiments are polynucleotides encoding EBI 3 variants comprising the amino acid sequence of EBI 3 polypeptide of Table 2 (SEQ ID NO:2) in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acid residues are substituted, deleted or added, in any combination.

[0037] The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 80%, and preferably at least 90%, and more preferably at least 95%, yet even more preferably 97-99% identity between the sequences.

[0038] Polynucleotides of the invention, which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1 or a fragment thereof, may be used as hybridization probes for cDNA and genomic DNA, to isolate full-length cDNAs and genomic clones encoding EBI 3 and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to the EBI 3 gene. Such hybridization techniques are known to those of skill in the art. Typically these nucleotide sequences are 80% identical, preferably 90% identical, more preferably 95% identical to that of the referent. The probes generally will comprise at least 15 nucleotides. Preferably, such probes will have at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will range between 30 and 50 nucleotides.

[0039] In one embodiment, to obtain a polynucleotide encoding EBI 3 polypeptide, including homologs and orthologs from species other than human, comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the SEQ ID NO: 1 or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Thus in another aspect, EBI 3 polynucleotides of the present invention further include a nucleotide sequence comprising a nucleotide sequence that hybridize under stringent condition to a nucleotide sequence having SEQ ID NO: 1 or a fragment thereof. Also included with EBI 3 polypeptides are polypeptide comprising amino acid sequence encoded by nucleotide sequence obtained by the above hybridization condition. Such hybridization techniques are well known to those of skill in the art. Stringent hybridization conditions are as defined above or, alternatively, conditions under overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

[0040] The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments and diagnostics to animal and human disease.

Vectors, Host Cells, Expression

[0041] The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, and host cells which are genetically engineered with vectors of the invention and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

[0042] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY* (1986) and Sambrook *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) such as calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

[0043] Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

[0044] A great variety of expression systems can be used. Such systems include, among others, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof such as those derived from plasmid and bacteriophage genetic

elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL (supra)*.

[0045] For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the desired polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0046] If the EBI 3 polypeptide is to be expressed for use in screening assays, generally, it is preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If EBI 3 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide; if produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

[0047] EBI 3 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Diagnostic Assays

[0048] This invention also relates to the use of EBI 3 polynucleotides for use as diagnostic reagents. Detection of a mutated form of EBI 3 gene associated with a dysfunction will provide a diagnostic tool that can add to or define a diagnosis of a disease or susceptibility to a disease which results from under-expression, over-expression or altered expression of EBI 3. Individuals carrying mutations in the EBI 3 gene may be detected at the DNA level by a variety of techniques.

[0049] Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled EBI 3 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing. See, e.g., Myers *et al.*, *Science* (1985) 230:1242. Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method. See Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401. In another embodiment, an array of oligonucleotide probes comprising EBI 3 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability. (See for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

[0050] The diagnostic assays offer a process for diagnosing or determining a susceptibility to infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome through detection of mutation in the EBI 3 gene by the methods described.

[0051] In addition, infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, can be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of EBI 3 polypeptide or EBI 3 mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as an EBI 3, in a sample derived from a host are well-known to

those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

[0052] Thus in another aspect, the present invention relates to a diagnostic kit for a disease or susceptibility to a disease, particularly infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, which comprises:

- (a) a EBI 3 polynucleotide, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a EBI 3 polypeptide, preferably the polypeptide of SEQ ID NO: 2, or a fragment thereof; or
- (d) an antibody to a EBI 3 polypeptide, preferably to the polypeptide of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Chromosome Assays

[0053] The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

Antibodies

[0054] The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them can also be used as immunogens to produce antibodies immunospecific for the EBI 3 polypeptides. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

[0055] Antibodies generated against the EBI 3 polypeptides can be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp. 77-96, Alan R. Liss, Inc., 1985).

[0056] Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms including other mammals, may be used to express humanized antibodies.

[0057] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

[0058] Antibodies against EBI 3 polypeptides may also be employed to treat infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others.

Vaccines

[0059] Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with EBI 3 polypeptide, or a fragment thereof, adequate to produce antibody and/or T cell immune response to protect said animal from infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, among others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering EBI 3 polypeptide via a vector directing expression of EBI 3 polynucleotide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

[0060] Further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a EBI 3 polypeptide wherein the composition comprises a EBI 3 polypeptide or EBI 3 gene. The vaccine formulation may further comprise a suitable carrier. Since EBI 3 polypeptide may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc. injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Screening Assays

[0061] The EBI 3 polypeptide of the present invention may be employed in a screening process for compounds which bind the receptor and which activate (agonists) or inhibit activation of (antagonists) the receptor polypeptide of the present invention. Thus, polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See Coligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991).

[0062] EBI 3 polypeptides are responsible for many biological functions, including many pathologies. Accordingly, it is desirable to find compounds and drugs which stimulate EBI 3 on the one hand and which can inhibit the function of EBI 3 on the other hand. In general, agonists are employed for therapeutic and prophylactic purposes for such conditions as infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome. Antagonists may be employed for a variety of therapeutic and prophylactic purposes for such conditions as infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV-1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome.

[0063] In general, such screening procedures involve producing appropriate cells which express the receptor polypeptide of the present invention on the surface thereof. Such cells include cells from mammals, yeast, *Drosophila* or *E. coli*. Cells expressing the receptor (or cell membrane containing the expressed receptor) are then contacted with a test compound to observe binding, or stimulation or inhibition of a functional response.

[0064] One screening technique includes the use of cells which express receptor of this invention (for example, transfected CHO cells) in a system which measures extracellular pH or intracellular calcium changes caused by receptor activation. In this technique, compounds may be contacted with cells expressing the receptor polypeptide of the present invention. A second messenger response, e.g., signal transduction, pH changes, or changes in calcium level, is then measured to determine whether the potential compound activates or inhibits the receptor.

[0065] Another method involves screening for receptor inhibitors by determining inhibition or stimulation of receptor-mediated cAMP and/or adenylate cyclase accumulation. Such a method involves transfecting a eukaryotic cell with the receptor of this invention to express the receptor on the cell surface. The cell is then exposed to potential antagonists in the presence of the receptor of this invention. The amount of cAMP accumulation is then measured. If the potential antagonist binds the receptor, and thus inhibits receptor binding, the levels of receptor-mediated cAMP, or adenylate cyclase, activity will be reduced or increased.

[0066] Another method for detecting agonists or antagonists for the receptor of the present invention is the yeast based technology as described in U.S. Patent 5,482,835.

[0067] The assays may simply test binding of a candidate compound wherein adherence to the cells bearing the receptor is detected by means of a label directly or indirectly associated with the candidate compound or in an assay involving competition with a labeled competitor. Further, these assays may test whether the candidate compound results in a signal generated by activation of the receptor, using detection systems appropriate to the cells bearing the receptor at their surfaces. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed.

[0068] Further, the assays may simply comprise the steps of mixing a candidate compound with a solution containing a EBI 3 polypeptide to form a mixture, measuring EBI 3 activity in the mixture, and comparing the EBI 3 activity of the mixture to a standard.

[0069] The EBI 3 cDNA, protein and antibodies to the protein may also be used to configure assays for detecting the effect of added compounds on the production of EBI 3 mRNA and protein in cells. For example, an ELISA may be constructed for measuring secreted or cell associated levels of EBI 3 protein using monoclonal and polyclonal antibodies by standard methods known in the art, and this can be used to discover agents which may inhibit or enhance the production of EBI 3 (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues. Standard methods for conducting screening assays are well understood in the art.

[0070] Examples of potential EBI 3 antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligand of the EBI 3, e.g., a fragment of the ligand, or small molecules which bind to the receptor but do not elicit a response, so that the activity of the receptor is prevented.

[0071] Thus in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for EBI 3 polypeptides; or compounds which decrease or enhance the production of EBI 3 polypeptides, which comprises:

- (a) a EBI 3 polypeptide, preferably that of SEQ ID NO:2;
- (b) a recombinant cell expressing a EBI 3 polypeptide, preferably that of SEQ ID NO:2;
- (c) a cell membrane expressing a EBI 3 polypeptide; preferably that of SEQ ID NO: 2; or
- (d) antibody to a EBI 3 polypeptide, preferably that of SEQ ID NO: 2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

Prophylactic and Therapeutic Methods

[0072] This invention provides methods of treating abnormal conditions such as, infections such as bacterial, fungal, protozoan and viral infections, particularly infections caused by HIV- 1 or HIV-2; pain; cancers; anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; ulcers; asthma; allergies; benign prostatic hypertrophy; and psychotic and neurological disorders, including anxiety, schizophrenia, manic depression, delirium, dementia, severe mental retardation and dyskinesias, such as Huntington's disease or Gilles de la Tourette's syndrome, related to both an excess of and insufficient amounts of EBI 3 activity.

[0073] If the activity of EBI 3 is in excess, several approaches are available. One approach comprises administering to a subject an inhibitor compound (antagonist) as hereinabove described along with a pharmaceutically acceptable carrier in an amount effective to inhibit activation by blocking binding of ligands to the EBI 3, or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of EBI 3 polypeptides still capable of binding the ligand in competition with endogenous EBI 3 may be administered. Typical embodiments of such competitors comprise fragments of the EBI 3 polypeptide.

[0074] In still another approach, expression of the gene encoding endogenous EBI 3 can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered. See, for example, O'Connor, *J Neurochem* (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Alternatively, oligonucleotides which form triple helices with the gene can be supplied. See, for example, Lee *et al.*, *Nucleic Acids Res* (1979) 6:3073; Cooney *et al.*, *Science* (1988) 241:456; Dervan *et al.*, *Science* (1991) 251:1360. These oligomers can be administered *per se* or

the relevant oligomers can be expressed *in vivo*.

[0075] For treating abnormal conditions related to an under-expression of EBI 3 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates EBI 3, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of EBI 3 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of EBI 3 polypeptides in combination with a suitable pharmaceutical carrier.

Formulation and Administration

[0076] Peptides, such as the soluble form of EBI 3 polypeptides, and agonists and antagonist peptides or small molecules, may be formulated in combination with a suitable pharmaceutical carrier. Such formulations comprise a therapeutically effective amount of the polypeptide or compound, and a pharmaceutically acceptable carrier or excipient. Such carriers include but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. Formulation should suit the mode of administration, and is well within the skill of the art. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

[0077] Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0078] Preferred forms of systemic administration of the pharmaceutical compositions include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if properly formulated in enteric or encapsulated formulations, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels and the like.

[0079] The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 $\mu\text{g/kg}$ of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

[0080] Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

Examples

[0081] The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples illustrate, but do not limit the invention.

Example 1:

[0082] A RT-PCR was performed on human spleen mRNA using a pair of primers specific for human EBI 1 gene. The products were subcloned into pCR2. 1 vector (Invitrogene). Multiple clones were selected for sequencing. One of the clones, EBI 3, was identified as a splicing variant of the EBI 1, which has the internal in frame deletion, missing 17 amino acids.

Example 2: Mammalian Cell Expression

[0083] The receptors of the present invention are expressed in either human embryonic kidney 293 (HEK293) cells or adherent dhfr CHO cells. To maximize receptor expression, typically all 5' and 3' untranslated regions (UTRs) are removed from the receptor cDNA prior to insertion into a pCDN or pCDNA3 vector. The cells are transfected with individual receptor cDNAs by lipofectin and selected in the presence of 400 mg/ml G418. After 3 weeks of selection, individual clones are picked and expanded for further analysis. HEK293 or CHO cells transfected with the vector alone serve as negative controls. To isolate cell lines stably expressing the individual receptors, about 24 clones are typically selected and analyzed by Northern blot analysis. Receptor mRNAs are generally detectable in about 50% of the G418-resistant clones analyzed.

Example 3 Ligand bank for binding and functional assays.

[0084] A bank of over 200 putative receptor ligands has been assembled for screening. The bank comprises: transmitters, hormones and chemokines known to act via a human seven transmembrane (7TM) receptor; naturally occurring compounds which may be putative agonists for a human 7TM receptor, non-mammalian, biologically active peptides for which a mammalian counterpart has not yet been identified; and compounds not found in nature, but which activate 7TM receptors with unknown natural ligands. This bank is used to initially screen the receptor for known ligands, using both functional (i.e. calcium, cAMP, microphysiometer, oocyte electrophysiology, etc, see below) as well as binding assays.

Example 4: Ligand Binding Assays

[0085] Ligand binding assays provide a direct method for ascertaining receptor pharmacology and are adaptable to a high throughput format. The purified ligand for a receptor is radiolabeled to high specific activity (50-2000 Ci/mmol) for binding studies. A determination is then made that the process of radiolabeling does not diminish the activity of the ligand towards its receptor. Assay conditions for buffers, ions, pH and other modulators such as nucleotides are optimized to establish a workable signal to noise ratio for both membrane and whole cell receptor sources. For these assays, specific receptor binding is defined as total associated radioactivity minus the radioactivity measured in the presence of an excess of unlabeled competing ligand. Where possible, more than one competing ligand is used to define residual nonspecific binding.

Example 5: Functional Assay in Xenopus Oocytes

[0086] Capped RNA transcripts from linearized plasmid templates encoding the receptor cDNAs of the invention are synthesized in vitro with RNA polymerases in accordance with standard procedures. In vitro transcripts are suspended in water at a final concentration of 0.2 mg/ml. Ovarian lobes are removed from adult female toads, Stage V defolliculated oocytes are obtained, and RNA transcripts (10 ng/oocyte) are injected in a 50 nl bolus using a microinjection apparatus. Two electrode voltage clamps are used to measure the currents from individual Xenopus oocytes in response to agonist exposure. Recordings are made in Ca^{2+} free Barth's medium at room temperature. The Xenopus system can be used to screen known ligands and tissue/cell extracts for activating ligands.

Example 6: Microphysiometric Assays

[0087] Activation of a wide variety of secondary messenger systems results in extrusion of small amounts of acid from a cell. The acid formed is largely as a result of the increased metabolic activity required to fuel the intracellular signaling process. The pH changes in the media surrounding the cell are very small but are detectable by the CYTOSENSOR microphysiometer (Molecular Devices Ltd., Menlo Park, CA). The CYTOSENSOR is thus capable of detecting the activation of a receptor which is coupled to an energy utilizing intracellular signaling pathway such as the G-protein coupled receptor of the present invention.

Example 7: Extract/Cell Supernatant Screening

[0088] A large number of mammalian receptors exist for which there remains, as yet, no cognate activating ligand (agonist). Thus, active ligands for these receptors may not be included within the ligands banks as identified to date. Accordingly, the 7TM receptor of the invention is also functionally screened (using calcium, cAMP, microphysiometer, oocyte electrophysiology, etc., functional screens) against tissue extracts to identify natural ligands. Extracts that produce positive functional responses can be sequentially subfractionated until an activating ligand is isolated identified.

Example 8: Calcium and cAMP Functional Assays

5 [0089] 7TM receptors which are expressed in HEK 293 cells have been shown to be coupled functionally to activation of PLC and calcium mobilization and/or cAMP stimulation or inhibition. Basal calcium levels in the HEK 293 cells in receptor-transfected or vector control cells were observed to be in the normal, 100 nM to 200 nM, range. HEK 293 cells expressing recombinant receptors are loaded with fura 2 and in a single > 150 selected ligands or tissue/cell extracts are evaluated for agonist induced calcium mobilization. Similarly, HEK 293 cells expressing recombinant receptors are evaluated for the stimulation or inhibition of cAMP production using standard cAMP quantitation assays. Agonists presenting a calcium transient or cAMP fluctuation are tested in vector control cells to determine if the response is unique to the transfected cells expressing receptor.

10 [0090] All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5

(1) GENERAL INFORMATION

10

(i) APPLICANT: SmithKline Beecham Corporation

15

(ii) TITLE OF THE INVENTION: A Novel Splicing Variant of
the Epstein-Barr Virus-Induced G-Protein Coupled
Receptor

20

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SmithKline Beecham, Corporate Intellectual
Property

25

(B) STREET: Two New Horizons Court

(C) CITY: Brentford

(D) COUNTY: Middlesex

(E) COUNTRY: United Kingdom

30

(F) POST CODE: TW8 9EP

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

35

(B) COMPUTER: IBM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ for Windows Version 2.0

40

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: TO BE ASSIGNED

(B) FILING DATE: 29-JUL-1997

45

(C) CLASSIFICATION: UNKNOWN

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER:

50

(B) FILING DATE:

55

(viii) ATTORNEY/AGENT INFORMATION:

- 5 (A) NAME: CONNELL, Anthony Christopher
 (B) GENERAL AUTHORISATION NUMBER: 5630
 (C) REFERENCE/DOCKET NUMBER: GP-70177

(ix) TELECOMMUNICATION INFORMATION:

- 10 (A) TELEPHONE: +44 127 964 4395
 (B) TELEFAX: +44 181 975 6294
 (C) TELEX:

15

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 1094 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30 ACCATGGATT TGGGTAAACC AATGAAAAGC GTGCTGGTGG TGGCTCTCCT TGTCATTTTC 60
 CAGGTATGCC TGTGTCAAGA TGAGGTCACG GACGATTACA TCGGAGACAA CACCACAGTG 120
 GACTACACTT TGTTGAGTC TTTGTGCTCC AAGAAGGACG TCGGAACTT TAAAGCCTGG 180
 TTCCTCCCTA TCATGTACTC CATCATTTGT TTCGTGGGCC TACTGGGCAA TGGGCTGGTC 240
 35 GTGTTGACCT ATATCTATTT CAAGAGGCTC AAGACCATGA CCGATACCTA CCTGCTCAAC 300
 CTGGCGGTGG CAGACATCCT CTTCTCCTG ACCCTTCCCT TCTGGGCCTA CAGCGCGGCC 360
 AAGTCCTGGG TCTTCAGTGG CATGCTCCTA CTTCTTTGCA TCAGCATTGA CCGCTACGTG 420
 GCCATCGTCC AGGCTGTCTC AGCTCACCGC CACCGTGCCC GCGTCCTTCT CATCAGCAAG 480
 40 CTGTCCTGTG TGGGCATCTG GATACTAGGC ACAGTGCTCT CCATCCCAGA GCTCCTGTAC 540
 AGTGACCTCC AGAGGAGCAG CAGTGAGCAA GCGATGCGAT GCTCTCTCAT CACAGAGCAT 600
 GTGGAGGCCT TTATCACCAT CCAGGTGGCC CAGATGGTGA TCGGCTTTCT GGTCCCCCTG 660
 CTGGCCATGA GCTTCTGTGA CTTGTGTCATC ATCCGCACCC TGCTCCAGGC ACGCAACTTT 720
 45 GAGCGCAACA AGGCCATCAA GGTGATCATC GCTGTGGTCG TGGTCTTCAT AGTCTTCCAG 780
 CTGCCCTACA ATGGGGTGGT CCTGGCCCAG ACGGTGGCCA ACTTCAACAT CACCAGTAGC 840
 ACCTGTGAGC TCAGTAAGCA ACTCAACATC GCCTACGACG TCACCTACAG CCTGGCCTGC 900
 GTCCGCTGCT GCGTCAACCC TTTCTTGAC GCCTTCATCG GCGTCAAGTT CCGCAACGAT 960
 50 CTCTTCAAGC TCTTCAAGGA CCTGGGCTGC CTCAGCCAGG AGCAGCTCCG GCAGTGGTCT 1020
 TCCTGTCGGC ACATCCGGCG CTCCTCCATG AGTGTGGAGG CCGAGACCAC CACCACCTTC 1080
 TCCCCATAAT CTAG 1094

55

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 361 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Asp Leu Gly Lys Pro Met Lys Ser Val Leu Val Val Ala Leu Leu
 1 5 10 15
 Val Ile Phe Gln Val Cys Leu Cys Gln Asp Glu Val Thr Asp Asp Tyr
 20 25 30
 Ile Gly Asp Asn Thr Thr Val Asp Tyr Thr Leu Phe Glu Ser Leu Cys
 35 40 45
 Ser Lys Lys Asp Val Arg Asn Phe Lys Ala Trp Phe Leu Pro Ile Met
 50 55 60
 Tyr Ser Ile Ile Cys Phe Val Gly Leu Leu Gly Asn Gly Leu Val Val
 65 70 75 80
 Leu Thr Tyr Ile Tyr Phe Lys Arg Leu Lys Thr Met Thr Asp Thr Tyr
 85 90 95
 Leu Leu Asn Leu Ala Val Ala Asp Ile Leu Phe Leu Leu Thr Leu Pro
 100 105 110
 Phe Trp Ala Tyr Ser Ala Ala Lys Ser Trp Val Phe Ser Gly Met Leu
 115 120 125
 Leu Leu Leu Cys Ile Ser Ile Asp Arg Tyr Val Ala Ile Val Gln Ala
 130 135 140
 Val Ser Ala His Arg His Arg Ala Arg Val Leu Leu Ile Ser Lys Leu
 145 150 155 160
 Ser Cys Val Gly Ile Trp Ile Leu Ala Thr Val Leu Ser Ile Pro Glu
 165 170 175
 Leu Leu Tyr Ser Asp Leu Gln Arg Ser Ser Ser Glu Gln Ala Met Arg
 180 185 190
 Cys Ser Leu Ile Thr Glu His Val Glu Ala Phe Ile Thr Ile Gln Val
 195 200 205
 Ala Gln Met Val Ile Gly Phe Leu Val Pro Leu Leu Ala Met Ser Phe
 210 215 220
 Cys Tyr Leu Val Ile Ile Arg Thr Leu Leu Gln Ala Arg Asn Phe Glu
 225 230 235 240
 Arg Asn Lys Ala Ile Lys Val Ile Ile Ala Val Val Val Val Phe Ile

25

30

1. An isolated polynucleotide comprising a nucleotide sequence that has at least 95% identity over its entire length to a nucleotide sequence encoding the EBI 3 polypeptide of SEQ ID NO:2; or a nucleotide sequence complementary to said isolated polynucleotide.
2. The polynucleotide of claim 1 wherein said polynucleotide comprises the nucleotide sequence contained in SEQ ID NO:1 encoding the EBI 3 polypeptide of SEQ ID NO:2.
3. The polynucleotide of claim 1 wherein said polynucleotide comprises a nucleotide sequence that is at least 95% identical to that of SEQ ID NO: 1 over its entire length.
4. The polynucleotide of claim 3 which is polynucleotide of SEQ ID NO: 1.
5. The polynucleotide of claim 1 which is DNA or RNA.
6. A DNA or RNA molecule comprising an expression system, wherein said expression system is capable of producing a EBI 3 polypeptide comprising an amino acid sequence, which has at least 95% identity with the polypeptide of SEQ ID NO:2 when said expression system is present in a compatible host cell.
7. A host cell comprising the expression system of claim 6.
8. A process for producing a EBI 3 polypeptide comprising culturing a host of claim 7 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.
9. A process for producing a cell which produces a EBI 3 polypeptide thereof comprising transforming or transfecting a host cell with the expression system of claim 6 such that the host cell, under appropriate culture conditions, produces a EBI 3 polypeptide.
10. A EBI 3 polypeptide comprising an amino acid sequence which is at least 95% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.
11. The polypeptide of claim 10 which comprises the amino acid sequence of SEQ ID NO:2.

12. An antibody immunospecific for the EBI 3 polypeptide of claim 10.

13. A method for the treatment of a subject in need of enhanced activity or expression of EBI 3 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an agonist to said receptor; and/or
- (b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence that has at least 95% identity to a nucleotide sequence encoding the EBI 3 polypeptide of SEQ ID NO:2 over its entire length; or a nucleotide sequence complementary to said nucleotide sequence in a form so as to effect production of said polypeptide activity *in vivo*.

14. A method for the treatment of a subject having need to inhibit activity or expression of EBI 3 polypeptide of claim 10 comprising:

- (a) administering to the subject a therapeutically effective amount of an antagonist to said receptor; and/or
- (b) administering to the subject a nucleic acid molecule that inhibits the expression of the nucleotide sequence encoding said receptor; and/or
- (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said receptor for its ligand.

15. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of EBI 3 polypeptide of claim 10 in a subject comprising:

- (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said EBI 3 polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of the EBI 3 polypeptide expression in a sample derived from said subject.

16. A method for identifying agonists to EBI 3 polypeptide of claim 10 comprising:

- (a) contacting a cell which produces a EBI 3 polypeptide with a candidate compound; and
- (b) determining whether the candidate compound effects a signal generated by activation of the EBI 3 polypeptide.

17. An agonist identified by the method of claim 16.

18. A method for identifying antagonists to EBI 3 polypeptide of claim 10 comprising:

- (a) contacting a cell which produces a EBI 3 polypeptide with an agonist; and
- (b) determining whether the signal generated by said agonist is diminished in the presence of a candidate compound.

19. An antagonist identified by the method of claim 18.

20. A recombinant host cell produced by a method of Claim 9 or a membrane thereof expressing a EBI 3 polypeptide.